

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Treflan* TR-10*

EPA Reg. No. 62719-131

24(c) Special Local Need Registration SLN OR-900019
(For Distribution and Use Only in the State of Oregon)

In Oregon this label is valid until December 31, 2007 or until otherwise amended, withdrawn, canceled, or suspended.

Dodder Control in Established Alfalfa Grown for Seed

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan* TR-10* herbicide before applying. Carefully follow use precautions and applicable use directions.
- Use of Treflan TR-10 according to this supplemental labeling is subject to all precautions and limitations imposed by the EPA-registered label affixed to the product container.

Active Ingredient:

trifluralin: α,α,α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine.....	10%
Inert Ingredients	90%
Total	100%

For use only on alfalfa grown for seed fields under the control of members of the Oregon Alfalfa Seed Growers Association, Gardena Alfalfa Seed Growers Association, Inc., and Josephine Growers' Cooperative Association, and located in Malheur, Umatilla, Morrow, Josephine or Jackson Counties of Oregon.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as butyl rubber ≥ 11 mils; or natural rubber ≥ 14 mils; or neoprene rubber ≥ 14 mils; or nitrile rubber ≥ 14 mils
- Shoes plus socks

General Information

Treflan* TR-10* is a selective preemergence herbicide for control of many annual grasses and broadleaf weeds. Treflan TR-10 controls susceptible weeds as they germinate. Treflan TR-10 does not control established weeds.

General Use Precautions

Applied according to directions and under normal growing conditions, Treflan TR-10 will not harm the treated crop. Over-application may result in crop injury or rotational crop damage from soil residue. Uneven application or improper incorporation of Treflan TR-10 can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Treflan TR-10. Under these conditions, delayed crop development or reduced yields may result.

Do not apply Treflan TR-10 to soils that are wet or are subject to prolonged periods of flooding as poor weed control may result.

Rotation Crop Restrictions

Sugar Beets, Red Beets and Spinach: In Oregon, sugar beets, red beets or spinach should not be planted for 12 months after a spring application or 14 months after a fall application of Treflan TR-10. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application of Treflan TR-10. Moldboard plowing to a depth of 12 inches prior to planting these crops will reduce the possibility of crop injury.

Proso Millet, Corn, Sorghum (Milo), Oats and Annual or Perennial Grass Crops or Grass Mixtures: In Oregon, unless crop injury is acceptable, proso millet, corn, sorghum (milo), oats and annual or perennial grass crops or grass mixtures should not be planted for 12 months after a spring application or 14 months after a fall application of Treflan TR-10 to avoid the possibility of crop injury. If land has not been irrigated, these crops should not be planted for 18 months after a spring application or 20 months after a fall application. Moldboard plowing to a depth of 12 inches before planting these crops will reduce the possibility of crop injury.

Other Crops: Vegetable crops other than those to which Treflan TR-10 may be applied as a preplant soil incorporated treatment, should not be planted within 5 months after an application of Treflan TR-10.

Dodder Control in Established Alfalfa Grown for Seed

Treflan TR-10 is registered for general weed control in established alfalfa at up to 20 lb per acre. Under this Special Local Need Registration Treflan TR-10 may be surface applied to established alfalfa grown for seed only, at 30 to 40 lb per acre. When Treflan TR-10 is applied at more than 20 lb per acre, no portion of the treated field, including seed, seed screenings, hay, forage or stubble, may be used for human or animal feed (see Use Restrictions below).

Application Directions: Apply at a broadcast rate of 30 to 40 lb per acre in late winter or spring. Application and activation must occur before dodder germination. For best results Treflan TR-10 should be activated within 3 days after application. When surface applied, Treflan TR-10 may be activated by rainfall, overhead sprinkler irrigation, flood irrigation or furrow irrigation. If activated using furrow irrigation, the surface of beds between furrows should be thoroughly wetted.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling (using greater than 20 lb per acre of Treflan TR-10) is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use on Crops Grown for Seed. If the applicator of this pesticide is not the producer, the applicator should provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, should provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed. This restriction pertains to, but is not limited to, green chop, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies.”

Use Precautions:

- Where alfalfa is to be rotated to another crop in the season following application of Treflan TR-10, plant only those crops to which Treflan may be applied as a preplant incorporated treatment or crop injury may occur. Read and carefully follow all rotational crop restrictions on the label for Treflan TR-10. The potential for rotational crop injury may be reduced by moldboard plowing to a depth of 12 inches prior to planting.
- Do not apply to seedling alfalfa (before the 6 true leaf stage).

*Trademark of Dow AgroSciences LLC
R048-001
Approved: 01/14/04
Replaces D06-048-006.

Revisions: (1) Changed PPE in Ag Use Requirements box to match Sec 3 label; (2) revised and reformatted according to Oregon 24(c) labeling policy.