

RESTRICTED USE PESTICIDE

DUE TO AQUATIC TOXICITY, AVIAN HAZARD, ACUTE ORAL TOXICITY AND EYE CORROSIVITY HAZARD
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING FOR SPECIAL LOCAL NEED IN ARIZONA FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARIZONA

Supracide® 2E

EPA Reg No. 10163-236 / EPA SLN NO. AZ-000005

For control of certain insects in alfalfa grown for seed

ACTIVE INGREDIENT:	% BY WT.
Methidathion: O,O-dimethyl phosphorodithioate, S-ester with 4-(mercaptomethyl)-2-methoxy- Δ^2 -1,3,4-thiadiazolin-5-one	24.4%
OTHER INGREDIENTS	75.6%
TOTAL	100.0%

Supracide 2E contains 2 lbs. active ingredient per gallon.
Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN
 **DANGER/POISON** 
PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered Supracide 2E label.

This label must be in the possession of the user at the time of pesticide application.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

General Information

Alfalfa (Grown for Seed)

Apply at 4 pints of product per acre in a minimum of 10 gals. of water per acre with ground equipment or a minimum of 5 gals. of water per acre by air. Make foliar applications after spring cutback for control of Lygus, leafhoppers, and weevil. A maximum of one application of Supracide per season is allowed.

Note: To avoid injury to pollinating insects, do not apply during bloom if bees are in field area. Do not apply or allow drift to blooming crops or weeds when bees are actively foraging in the treatment area.

Precautions: Do not feed or graze treated growth of alfalfa forage or fodder. Do not cut treated growth of alfalfa for hay or forage. Seeds from treated plants may not be used for sprouts. No portion

of the treated field, including seed, seed screenings, hay or forage may be used for human or animal consumption.

Producers of alfalfa seed who use this product, or cause the product to be used on fields they operate, are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of the labeling is required to be provided to the conditioner by the producer. Processed seed must be labeled "Not for human or animal consumption" at the processing plant. The processor must dispose of all seed screenings in such a way that they cannot be distributed or used for food or feed.

Notes: Field reentry after application is 72 hours.

Chemigation: Do not apply this product through any type of irrigation system.

24(c) registrant:
Gowan Company
PO Box 5569
Yuma, AZ 85366-5569

EPA SLN No.:

AZ-000005

Supracide[®] is a trademark of Syngenta Crop Protection Inc.