

**RESTRICTED USE PESTICIDE**

DUE TO AQUATIC TOXICITY, AVIAN HAZARD, ACUTE ORAL TOXICITY AND EYE CORROSIVITY HAZARD  
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABELING FOR SPECIAL LOCAL NEED IN MONTANA  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA**

**NOTE:** A NONFOOD USE AGREEMENT MUST BE SIGNED WITH THE MONTANA DEPARTMENT OF AGRICULTURE PRIOR TO PURCHASE OF THIS PRODUCT

## Supracide® 2E

EPA REG. NO. 10163-236 / EPA SLN NO. MT-000008

### FOR CONTROL OF CERTAIN INSECTS IN ALFALFA GROWN FOR SEED

ACTIVE INGREDIENT:	% BY WT.
Methidathion: O,O-dimethyl phosphorodithioate, S-ester with 4-(mercaptomethyl)-2-methoxy- $\Delta^2$ -1,3,4-thiadiazolin-5-one	24.4%
OTHER INGREDIENTS	75.6%
<b>TOTAL</b>	<b>100.0%</b>

Supracide 2E contains 2 lbs. active ingredient per gallon.  
Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN**  
 **DANGER/POISON**   
**PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered Supracide 2E label.  
This label must be in the possession of the user at the time of pesticide application.

---

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.  
FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

---

#### General Information

##### Alfalfa (Grown for Seed)

Apply at the rates listed below in a minimum of 10 gals. of water per acre with ground equipment or a minimum of 2 gals. of water per acre by air. For alfalfa weevil and Egyptian alfalfa weevil control, make foliar applications to standing alfalfa when 20 - 30% of the growing tips show insect feeding damage. For other insects listed, apply when insects first appear. Use the lower rate for light infestations and the higher rate for moderate to heavy infestations and follow state IPM guidelines.

Note: To avoid injury to pollinating insects, do not apply during bloom. Do not apply or allow to drift to blooming crops or weeds when bees are actively foraging in the treatment area.

<b>Insects Controlled</b>	<b>Rates Per Acre</b>
Alfalfa Weevil	2-4 pints.
Aphids	2-4 pints.
Egyptian alfalfa weevil	2-4 pints.
Lygus	2-4 pints.
Leafhoppers	2-4 pints.

Precautions: Do not feed or graze current year's alfalfa forage or fodder. Do not cut current year's alfalfa for hay or forage. Seeds from treated fields may not be used for sprouts. No portion of the treated field, including seed, seed screenings, hay, forage, or stubble may be used for human or animal consumption.

Producers of alfalfa seed who use this product, or cause the product to be used on fields they operate, are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of the labeling is required to be provided to the conditioner by the producer. Processed seed must be labeled "Not for human or animal consumption" at the processing plant. The processor must dispose of all seed screenings in such a way that they cannot be distributed or used for food or feed.

Chemigation: Do not apply this product through any type of irrigation system.

24(c) registrant:  
Gowan Company  
PO Box 5569  
Yuma, AZ 85366-5569

EPA SLN No.:  
Effective Date:

MT-000008  
05-25-00

Supracide® is a registered trademark of Novartis