

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban*-4E

EPA Reg. No. 62719-220

EPA 24(c) Special Local Need Registration SLN NV-940002

For Distribution and Use Only in the States of Nevada

To Control Billbugs, Cutworms and Aphids Infesting Perennial Grass Seed Crops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban-4E before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban-4E according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban-4E.

Directions for Use

Use and Dosage Recommendations

To control billbugs and cutworms, apply Lorsban*-4E insecticide at a rate of 1 quart per acre as a broadcast spray using power-operated ground application. For best results, use 20 gallons of spray mixture per acre. For improved results, the thatch should be moist at the time of application and the insecticide washed into the thatch immediately after application. If sprinkler irrigation is available, apply 1/4 to 1/2 inch of water after application to help incorporate the material. Otherwise, apply to early morning dew-covered fields or during or immediately following a light rain. For Billbug control, apply in early April when adult billbugs are active. For cutworm control, apply in either spring or fall when larvae or larval damage first appears. Fall applications should follow sufficient irrigation and/or rainfall. Heavy carbon residue from field burning will prevent penetration of Lorsban-4E into the plant crown where the insects feed. Retreat as necessary to maintain control.

For aphid control, apply Lorsban-4E at a rate of 1 to 2 pints per acre as a broadcast spray using power-operated ground application equipment. For best results, use 20 gallons of spray mixture per acre. Treat before heading as colonies begin to form.

Consult your cooperative agricultural extension service for additional information concerning billbug, cutworm, and aphid control practices in your area.

Restrictions

- Apply only to perennial grass seed crops. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings or seed screenings to livestock or use hay for livestock bedding.
- This product is highly toxic to bees exposed to direct treatment or residues on treated crops or weeds. Protective information may be obtained from your cooperative agricultural extension service.

*Trademark of Dow AgroSciences LLC

D06-062-074

Approved: 04/26/94

Replaces 122-58-031N

Revisions:

Added Restricted Use Pesticide designation.